

Dissimilarity in the oxidative folding of onconase and ribonuclease A, two structural homologues

Robert F. Gahl, Mahesh Narayan¹, Guoqiang Xu² and Harold A. Scheraga³

Baker Laboratory of Chemistry and Chemical Biology, Cornell University, Ithaca, NY 14853-1301, ¹Present address: Department of Chemistry, University of Texas at El Paso, El Paso, TX 79968-0513 and ²Present address: Department of Pharmacology, Weill Medical College, Cornell University, New York, NY 10065, USA

³To whom correspondence should be addressed. E-mail: has5@cornell.edu

The oxidative folding of frog onconase (ONC), a member of the ribonuclease A family, was examined and shows markedly different behavior compared to its structural homologue bovine pancreatic ribonuclease A (RNase A) under similar conditions. Application of a reduction pulse (using a small amount of reduced dithiothreitol) during the oxidative regeneration of ONC indicated the survival of the native protein along with three other (structured) species, I₁, I₂ and I₃, with the rest of the unstructured species being converted to fully reduced protein. Mass spectrometry indicates that I₁ has two disulfide bonds, whereas I₂ and I₃ have three disulfide bonds each. A disulfide mapping method, based on cyanylation, was used to identify I₂ and I₃ as des-[30–75] and des-[19–68], respectively. On enzymatic digestion using trypsin, I₁ was identified as des-[19–68, 30–75]. Differences in the intermediates that are generated during the oxidative folding of the two structural homologues, RNase A and ONC, demonstrate that regenerative pathways are not necessarily influenced by tertiary structure. This indicates that the lack of a disulfide bond in ONC, analogous to the (65–72) disulfide bond in RNase A which plays an important role in its oxidative regeneration, does not adversely affect the oxidative folding of ONC.

Keywords: onconase/oxidative folding/RNase A/structured intermediate

Introduction

Onconase (ONC; registered trademark of Alfacell Corp., Bloomfield, NJ, USA), a frog ribonuclease found in the

Abbreviations: ONC, frog onconase (*Rana pipiens*); R-ONC, onconase with all its disulfide bonds reduced; RB-ONC, onconase with all its cysteines blocked with AEMTS and tagged with an $-\text{SCH}_2\text{CH}_2\text{NH}_3^+$ group; RNase A, bovine pancreatic ribonuclease A; AEMTS, 2-aminoethylmethylthiosulfonate; CDAP, 1-cyano-4-dimethylaminopyridinium tetrafluoroborate; DTT^{ox} and DTT^{red}, oxidized and reduced dithiothreitol, respectively; TCEP, tris(2-carboxyethyl)phosphine hydrochloride; GdnHCl, guanidine hydrochloride; Tris, tris(hydroxymethyl)aminomethane; HEPES, *N*-(2-hydroxyethyl)piperazine-*N'*-(2-ethanesulfonic acid); TFA, trifluoroacetic acid; SCX, strong cation exchange; RP, reversed phase; HPLC, high-performance liquid chromatography; des-[a–b, c–d], a species that contains all native disulfide bonds *except* for the a–b and c–d disulfide bonds; *n*S, an ensemble of disulfide-containing intermediates each having *n* disulfide bonds; *n*S*, a single species containing *n* disulfide bonds; N, native onconase; –itc, iminothiazolidinyl carboxyl residue; MALDI-TOF, matrix-assisted laser desorption/ionization time-of-flight.

oocytes and early embryos of the northern leopard frog (*Rana pipiens*) (Ardelt *et al.* 1991), has four disulfide bonds and is structurally homologous to bovine pancreatic ribonuclease A (RNase A) (Wlodawer *et al.* 1988; Mosimann *et al.* 1994; Leland and Raines 2001). The frog variant demonstrates cytostatic and cytotoxic characteristics toward certain types of tumor cells (Boix *et al.* 1996; Leland *et al.* 1998; Leland *et al.* 2000), can degrade tRNAs selectively (Saxena *et al.* 2002) and is currently in Phase IIIb clinical trials for the treatment of asbestos-related lung cancer (http://www.alfacell.com/clinical_trials/mm_trials.htm). Ongoing effort in various laboratories is devoted to understand and improve the biomedical potential of ONC, and has resulted in the discovery of several new and interesting properties of this macromolecule (Newton *et al.* 1998, 2001; Welker *et al.* 2007).

Research in our laboratory is devoted to determining the intramolecular interactions that govern the oxidative folding of multi-disulfide-containing proteins (Rothwarf and Scheraga 1993; Rothwarf *et al.* 1998a; Narayan *et al.* 2000; Wedemeyer *et al.* 2000; Welker *et al.* 2001a, 2001b, 2004). The ultimate aim was to be able to predict the structures of proteins from their amino acid sequences, which will aid in their applications in medicine and biotechnology. In this context, studies of the intermediates that accumulate during the folding pathway are useful in determining key interactions that allow for the formation and stabilization of those intermediates, and of the folded protein. Knowledge of such interactions is essential for the design of more potent mimics of the naturally occurring biomolecule, and is also required to improve the yields during the oxidative folding of recombinant native and mutant variants that are created for biotechnological and therapeutic purposes.

To understand the factors influencing the oxidative folding pathways of multi-disulfide-containing proteins, we have carried out in-depth studies of the regeneration of RNase A and its three-disulfide-containing mutants (Rothwarf and Scheraga 1993; Iwaoka *et al.* 1998; Rothwarf *et al.* 1998a; Xu and Scheraga 1998). Our work, together with research in other laboratories, has demonstrated that multi-disulfide-containing proteins regenerate by first forming unstructured disulfide-containing intermediates (Weissman and Kim 1991; van den Berg *et al.* 1999; Narayan *et al.* 2000; Chang and Li 2002). The rate determining steps in most of these cases involve the formation of a structured intermediate, or the native protein, either through reshuffling-coupled conformational folding reactions (such as the 3S→3S* step in RNase A (Rothwarf *et al.* 1998b) or through oxidation-coupled conformational folding reactions (such as the 2S→N [or 3S*] step in hirudin (Thannhauser *et al.* 1997) and in the three-disulfide-containing mutants of RNase A (Iwaoka *et al.* 1998; Xu and Scheraga 1998). The formation of the remainder of the disulfide bonds in a structured intermediate is more rapid in this kinetic mechanism and, thus, the

formation of such a structured intermediate plays an important role in the overall oxidative folding pathway.

In this study, we have examined the oxidative folding of ONC and the effects of any structured intermediates in the early stages of regeneration on the oxidative folding. We have also identified the disulfide-bonded residues in the oxidative folding intermediates. The results obtained are compared with the oxidative folding pathways in the regeneration of RNase A, for which a large amount of data already exists (Rothwarf and Scheraga 1993; Iwaoka *et al.* 1998; Rothwarf *et al.* 1998a, 1998b; Xu and Scheraga 1998; Narayan *et al.* 2000; Wedemeyer *et al.* 2000; Welker *et al.* 2001a, 2001b, 2004), and with the reductive unfolding pathways of RNase A and ONC (Narayan *et al.* 2004). Furthermore, we use these structural homologues to determine the influence of three-dimensional structure on the oxidative folding pathways of multi-disulfide-containing proteins. Finally, the impact of the absence of the (65–72) disulfide bond in the oxidative folding of ONC is discussed, given that this disulfide bond plays a major role in the regeneration of RNase A (Xu *et al.* 1996, 2004; Iwaoka and Juminaga *et al.* 1998; Rothwarf *et al.* 1998a, 1998b).

Materials and methods

Materials

WT-ONC cDNA in a pET11 expression vector was kindly provided by R.J. Youle (Boix *et al.* 1996). This cDNA was amplified by PCR and cloned to a pET22b(+) vector in-frame with the pelB signal sequence without the starting methionine residue. WT-ONC (Boix *et al.* 1996) was expressed in BL21 (DE3) cells and purified as described earlier for RNase A (Laity *et al.* 1993, 1997; Narayan *et al.* 2004), and the level of expression was comparable with that of an independent expression of WT-ONC from another wild-type plasmid (pONC) which was kindly provided by R.T. Raines (Leland *et al.* 1998). The conversion of the leading glutamine to pyro-glutamic acid was made from the uncyclized protein by dialyzing the folded protein against 100 mM phosphate buffer, pH 7.0, at room temperature for 2 days. The protein was purified by cation-exchange HPLC as described earlier (Laity *et al.* 1993, 1997), and lyophilized. The conversion of the leading glutamine to pyroglutamic acid was verified by MALDI-TOF mass spectrometry (Cornell University Life Sciences Core Laboratories Center Proteomics and Mass Spectrometry Facility).

AEMTS (>99% pure) was purchased from Anatrache and used without further purification. DTT^{ox}, DTT^{red}, CDAP, methylamine HCl and TCEP were obtained from Sigma and used without further purification. Trypsin Gold, mass spectroscopy grade, was purchased from Promega in vials of 100 µg and was dissolved in 50 mM acetic acid, pH 3, to a concentration of 1 mg/ml. Aliquots (10 µl) of this stock solution were stored frozen at –70°C. Such aliquots were not used after three freeze/thaw cycles. All other chemicals were of the highest grade commercially available.

Preparation of reduced ONC

Lyophilized ONC was dissolved in buffer (pH 8, 25 mM Tris–HCl, 3 mM EDTA) containing 6 M GdnHCl and 100 mM DTT^{red} at a concentration of 2 mg/ml and allowed

to incubate at room temperature for a period of 2 h under humidified argon. The reducing agent and salts were removed from the reduced protein by using a G25 size-exclusion column (50 mM acetic acid as the running buffer). Water and acetic acid were removed by lyophilization. Lyophilized, reduced-ONC (R-ONC) was stored in 2 mg/ml aliquots in a 3 mM acetic acid buffer (pH 3), frozen at –70°C, until further use.

Oxidative folding of R-ONC

Oxidative folding of ONC was initiated by introducing an aliquot of R-ONC into a solution containing 25 mM DTT^{ox} (final pH 8.0, 100 mM Tris, 3 mM EDTA, 25°C), which was previously purged of oxygen by humidified argon, to a final concentration of 0.2 mg/ml of protein. The reaction mixture was kept under humidified argon for the entire time required to populate the oxidative folding intermediates.

Aliquots of the reaction mixture were withdrawn periodically, and free thiols in each aliquot were blocked immediately with AEMTS. The blocking was performed by adding 100 µl of a blocking buffer (pH 8.7, 2M Tris, 20 mM EDTA) with a sufficient amount of AEMTS to achieve a 100-fold molar excess over the thiol content, and was allowed to proceed for 2 min, then stopped by adding glacial acetic acid; every sample was immediately desalted on a G25 column, with 50 mM acetic acid as a running buffer. The intermediate species populated during oxidative folding were separated from the AEMTS blocked mixture by HPLC using a TOSOH Biosciences TSKgelTM column SP-5PW 7.5 cm × 7.5 mm, analytical strong cation-exchange (SCX) column with a salt gradient (0–480 mM NaCl in 120 min). The mobile phase was 25 mM HEPES with 2 mM EDTA buffer, pH 8.0.

In a previous study, some aliquots were subjected to a reduction pulse (Rothwarf *et al.* 1998a) by the addition of 5 mM DTT^{red} for a period of 2 min at 25°C before being blocked with AEMTS. Three chromatographic intermediates, namely as I₁, I₂ and I₃, were observed (Welker *et al.* 2004); their survival during a reduction pulse implies that they are structured. These chromatographic intermediates are reproduced in Fig. 1. According to the mass spectrometric results in Table I, I₂ and I₃ each has three disulfide bonds, and I₁ has two disulfide bonds. Favorable conditions




Fig. 1. SCX-HPLC separation of the species involved in the oxidative folding of ONC at 25 mM DTT^{ox}, pH 8.0, 25°C, 0.35 mg/ml ONC, 50 min after the start of regeneration. The absorbance at 280 nm was monitored.

Table 1. Molecular weights of native ONC and its AEMTS-blocked intermediates, I₁, I₂ and I₃

Observed chromatographic peaks of Fig. 1	Experimental MW of ONC species in Fig. 1, blocked with AEMTS ^a	Theoretical MW of ONC species blocked with AEMTS	Number of disulfide bonds in each species in Fig. 1
N	11 820 ^b	11 819 ^{b,c}	4
I ₃	11 970	11 971	3
I ₂	11 971	11 971	3
I ₁	12 121	12 123	2
R	12 427	12 427	0

Each AEMTS adduct adds 76 Da to the molecular weight of the native protein.

^aMass measurements were made on a MALDI-TOF MS instrument.

^bN cannot be blocked with AEMTS.

^cArdelt *et al.* (1991).

were established to populate each of these intermediates in large enough quantities to determine their disulfide connectivities.

Populating the oxidative folding intermediates for disulfide mapping

To optimally populate I₁, oxidative folding was initiated and allowed to proceed for 15 min, before being blocked with AEMTS, and isolated from an SCX-HPLC separation, as described in 'Oxidative folding of R-ONC' section. After a fraction from the HPLC column containing I₁ was collected, the pH of the solution was reduced to 3, and the sample was frozen at -70°C until further use. NaCl and buffer salts were removed by RP-HPLC using a 25 cm \times 4.6 mm SULPELCO Discovery[®] BIO Wide Pore C18, 5 μm particle size column with a 0.1% TFA–water/0.1% TFA–acetonitrile mobile phase with a constant gradient of 0% acetonitrile to elute salts, followed by a constant gradient of 80% acetonitrile to elute I₁. After collecting AEMTS-blocked I₁ from RP-HPLC, the collected fraction was frozen by allowing the vessel to contact dry ice, prior to lyophilization (for removal of water, acetonitrile and TFA). The lyophilized powder was re-dissolved at pH 3 in 3 mM acetic acid buffer and frozen at -70°C until subjected to enzymatic digestion.

I₂ and I₃ were optimally populated by allowing the oxidative folding reaction to proceed for 60 min. Then, the reaction was stopped by reducing the pH to 3 by adding 100 μl of glacial acetic acid. They were then separated by RP-HPLC using a 25 cm \times 4.6 mm SULPELCO Discovery[®] BIO Wide Pore C18, 5 μm particle size column with 0.1% TFA–water/0.1% TFA–acetonitrile mobile phase and a gradient of 30–45% acetonitrile over 60 min. After I₂ and I₃ were collected in separate fractions, each fraction was frozen by allowing the vessel to contact dry ice and lyophilized to remove the TFA, water and acetonitrile. Unblocked I₂ and I₃ were stored frozen in solution either as a collected fraction from RP-HPLC or reconstituted from the lyophilized form in 3 mM acetic acid at -70°C .

Free cysteines in I₂ and I₃ are unblocked during their separation by RP-HPLC because the mapping method used (Wu and Watson 1997) requires that the cysteines, which eventually must be cyanylated, are free, i.e. not blocked with

AEMTS. To verify that the peaks separated by RP-HPLC were I₂ and I₃, each separate fraction containing a single moiety, corresponding to I₂ or I₃, was lyophilized, reconstituted in 3 mM acetic acid over ice for 15 min and blocked with AEMTS, as described in 'Oxidative folding of R-ONC' section. The separate fractions were injected individually into the SCX-HPLC column and analyzed by the same method described in 'Oxidative folding of R-ONC' section. Each of the fractions analyzed separately corresponded to the intermediates I₂ and I₃ according to the SCX-HPLC chromatogram shown in Fig. 1.

Populating the oxidative folding intermediates by reductive unfolding for disulfide mapping

In order to map each oxidative folding intermediate unambiguously, the connectivity of the disulfide bonds had to be verified. Since these intermediates show evidence of native-like structure, it seems that the intermediates have native disulfide bonds. In order to verify this, each intermediate was populated both by reductive unfolding of native ONC (N-ONC) and by oxidative folding of R-ONC, and the cysteines not involved in disulfide bonds were identified in both cases. The disulfide bond connectivity could be verified as native only when the two mass spectra of each intermediate are the same, whether it was populated from N-ONC or R-ONC. Co-elution tests were carried out to verify that an oxidative folding intermediate was also populated by reductive unfolding.

I₂ and I₃ were populated by reductive unfolding by introducing an aliquot of N-ONC into to a solution containing 6.6 mM TCEP (pH 3.1, 0.1 M citric acid, 6 M GdnHCl, 2 mM EDTA, 25 $^{\circ}\text{C}$) to a final concentration of 0.3 mg/ml for 1 h. In previous work (Narayan *et al.* 2004; Xu *et al.* 2004), only I₂ was observed upon reduction of native protein (25 $^{\circ}\text{C}$, pH 8), in the absence of denaturant. Therefore, to access the disulfide bond to be reduced to populate I₃, 6 M GdnHCl had to be added to expose all of the disulfide bonds in N-ONC. After 1 h, in the presence of GdnHCl, unblocked I₂ and I₃ could be separated by the same method and conditions used to separate the intermediates described in 'Populating the oxidative folding intermediates for disulfide mapping' section, i.e. a brief reduction time at a low concentration of TCEP. I₂ and I₃, populated by reductive unfolding and blocked with AEMTS, could thus be separated. After N-ONC was permitted to equilibrate in the reaction mixture for 1 h, the mixture was diluted by one-half with 100 mM Tris–HCl, 2 mM EDTA buffer, pH 8.3, and immediately added to a solution containing AEMTS, as described in 'Oxidative folding of R-ONC' section. After 2 min, the resulting mixture was desalted on a G25 column with 50 mM acetic acid as a running buffer. AEMTS-blocked I₂ and I₃ could be separated by SCX-HPLC, using the method described in 'Oxidative folding of R-ONC' section.

I₁ was also populated by first reducing N-ONC to I₂ and isolating unblocked I₂, then populating I₁ by reducing unblocked I₂. Unblocked I₂ was most efficiently populated from reductive unfolding by introducing an aliquot of N-ONC to a solution containing 10 mM DTT^{red} (pH 8.0, 100 mM Tris, 2 mM EDTA) to a final concentration of 1 mg/ml for 1 h. Glacial acetic acid was added to stop the reduction reaction. The solution was injected into an RP-HPLC column, and unblocked I₂ was separated and

collected by the method described in ‘Populating the oxidative folding intermediates for disulfide mapping’ section. The fraction containing unblocked I_2 was frozen by allowing the vessel to contact dry ice and lyophilized to remove water, acetonitrile and TFA. The lyophilized powder was re-dissolved in 3 mM acetic acid buffer, pH 3, at a concentration of 7.5 mg/ml of unblocked I_2 stock solution. A 20 μ l aliquot was removed from this stock solution and added to a reducing buffer containing 6.6 mM TCEP (pH=3.1, 6 M GdnHCl, 0.1 M Tris, 2 mM EDTA) to obtain a final protein concentration of 0.3 mg/ml, and sufficient amount of I_1 was produced after 45 min. In order to block I_1 with AEMTS, the mixture was treated in a similar manner as for the AEMTS-blocking of I_2 and I_3 and separated on an SCX-HPLC column, as described previously.

Wu–Watson cyanylation and base cleavage method employed for disulfide bond connectivity identification in I_2 and I_3

Location of free cysteines in I_2 and I_3 was achieved with a method developed by Wu and Watson (Wu and Watson 1997). This method makes use of a nucleophilic attack by a base on the carbonyl carbon of the residue located on the N-terminal side of a cyanylated cysteine, resulting in cleavage of the peptide bond (Fig. 2). Following the

base-induced cleavage, the free amino group of the cyanylated cysteine residue performs a nucleophilic attack on the cyanyl group and forms a five-member ring, abbreviated as –itc (Wu and Watson 1998). The cysteines not involved in a disulfide-bond in an oxidative folding intermediate are cyanylated by CDAP, and the free cyteines can be identified by the size of the fragments produced by the cleavage.

After fractions of unblocked I_2 and I_3 were separated and collected individually from RP-HPLC and lyophilized, the dried fractions were treated with a cyanylation buffer. For the cyanylation of I_2 , a 50 μ l mixture containing 0.1 M CDAP (pH 3, 0.1 M citric acid) was added to the lyophilized powder for 15 min at 25°C. For the cyanylation of I_3 , a 25 μ l mixture containing 0.2 M CDAP (pH 3, 0.1 M citric acid, 6 M GdnHCl) was added to the lyophilized powder for 15 min at 25°C. Cyanylation of both samples was quenched by freezing them. CDAP, GdnHCl and buffer salts were removed from each fraction by RP-HPLC, using the same method to remove salts described in ‘Populating the oxidative folding intermediates for disulfide mapping’ section. The collected intermediates were frozen by allowing the vessel to contact dry ice and lyophilized. The conditions for cyanylating I_2 and I_3 are different because of the different reactivity of the free cysteines in both intermediates; i.e. the free cysteines in I_3 are less reactive than those in I_2 and, as a




Fig. 2. The mapping method developed by Wu and Watson (1997) applied to a three-disulfide-bond-containing intermediate of ONC; this scheme is illustrated with I_2 as an example. The cysteines in ONC are numbered, and the native disulfide bonds are connected. The locations of the free cysteines are identified by the sizes of the fragments produced by the base cleavage.

result, a higher concentration of CDAP and the presence of 6 M GdnHCl are required for complete blocking. Reshuffling and incomplete cyanylation was observed if these two procedures were not adopted in cyanylating I_3 .

Fragmentation was carried out with a 50 μ l mixture containing 2 M methylamine (pH 11, 6 M GdnHCl, 0.1 M phosphate) at 2°C for 10 min. Before the fragmentation mixture was added to the dried cyanylated intermediates, the fragmentation mixture was equilibrated for 5 min at 2°C. After 10 min, the fragmentation was stopped by adding 40 μ l of formic acid, and the disulfide bonds in each intermediate were reduced by adding 800 μ l of 0.1 M TCEP (pH 3.1, 0.1 M citric acid) for 40 min at 37°C.

Trypsin digestion of AEMTS-blocked I_1

Tryptic digestion, without the addition of GdnHCl, was used to produce peptides, identifiable by mass spectroscopy, for the identification of I_1 . The peptides were compared to those produced from tryptic digestion of a completely reduced and AEMTS-blocked species of ONC, RB-ONC.

The conditions for enzymatic digestion were 37°C and pH 8.0 (0.1 M Tris-HCl, 2 mM EDTA). The concentration of the protein was 0.5 mg/ml, and a sufficient amount of trypsin was added to form an enzyme: I_1 mass ratio of 1:25. The digestion was stopped by adding formic acid after 1 h.

Obtaining mass spectra for each intermediate

Each sample was desalted using a Sep-Pak reversed-phase Solid-Phase Extraction procedure before being loaded onto a 4700 Proteomics Analyzer (Applied Biosystems) MALDI-TOF mass spectrometer. Peptides were analyzed in both linear and reflector modes.

Population of I_1 from I_3

I_1 was populated from I_3 first by reducing N-ONC to I_3 , isolating unblocked I_3 from RP-HPLC, and lyophilizing and storing the re-dissolved protein in 3 mM acetic acid, as discussed in 'Populating the oxidative folding intermediates by reductive unfolding for disulfide mapping' section. For the reduction of I_3 , this unblocked intermediate was added to a reducing buffer containing 0.1 M TCEP (pH 3.1, 0.1 M citric acid) to a final protein concentration of 0.1 mg/ml, and I_1 was produced after 30 min. I_3 and I_1 were separated by RP-HPLC according to 'Populating the oxidative folding intermediates for disulfide mapping' section.

Results

Oxidative folding of ONC

Figure 3A and B are typical HPLC chromatograms showing the oxidative regeneration of N-ONC from its reduced form as a function of time, 15 and 115 min, respectively (using 25 mM DTT^{ox}). During the course of oxidative folding, several intermediates are visible, besides the native (N) in Fig. 3A and B and fully-reduced (R) protein in Fig. 3A. After only 15 min, almost all of R and 1S are consumed. This is much different from the oxidative folding behavior of RNase A, in which R and 1S are in an identifiable pre-equilibrium in which they each constitute 10 and 20%, respectively, of the intermediate species over a number of hours under similar redox conditions (Rothwarf and




Fig. 3. Typical HPLC chromatograms showing the oxidative regeneration of ONC using DTT^{ox} (25 mM) at pH 8.0 and 25°C (100 mM Tris-HCl, 3 mM EDTA) at an R-ONC concentration of 0.2 mg/ml. (A) 15 min and (B) 115 min after the start of the regeneration. The samples were blocked with AEMTS before injection onto the HPLC, as described in 'Oxidative folding of R-ONC' section. The absorbance at 280 nm was monitored.

Scheraga 1993). This shows that strong interactions play a role early in the oxidative folding pathway of ONC, and drive formation of the early forming intermediate, I_1 . According to Fig. 3B, 40% of N is regenerated after 115 min. Compared with RNase A, this degree of folding in such a short amount of time is observed only with 200 mM DTT^{ox}, eight times the amount used for ONC. Another key difference is that the intermediates, I_1 , I_2 and I_3 , are present to a large extent, 10–30% over the course of the oxidative folding process, compared with RNase A. The stable intermediates in RNase A, des-[65–72] and des-[40–95] were populated to no more than 10% (Rothwarf *et al.* 1998b) under conditions much more oxidizing than the ones used for ONC in Fig. 3. Also, the reactivities of I_2 and I_3 along the oxidative folding pathway of ONC appear to be different from each other. In Fig. 3A, I_2 is produced to a larger extent than I_3 , whereas in Fig. 3B more I_2 is being turned over than I_3 . Further work (in progress) would determine how their different reactivities play a role in the oxidative folding mechanism.

Production of oxidative folding intermediates by reductive unfolding

The oxidative folding intermediates produced by the reductive unfolding of N and subsequently blocked with AEMTS are shown in Fig. 4. The chromatogram for this separation is superimposed on the chromatogram for separation of the AEMTS-blocked intermediates produced by oxidative folding. To show that the intermediates from reduction and oxidation are the same species, a co-elution test was carried out and the results (in the inset of Fig. 4) indicated that they are the same species. As pointed out in 'Populating the oxidative folding intermediates by reductive unfolding for disulfide mapping' section, only I_2 is observed in the reductive




Fig. 4. Superposition of the chromatograms of the oxidative folding intermediates produced by reduction of N-ONC and by oxidation of R-ONC. The conditions for reduction were: 25°C, 6.6 mM TCEP, pH 3, 6 M GdnHCl, 0.3 mg/ml N-ONC. The conditions for oxidation were: 25°C, 25 mM DTT^{ox}, pH 8.0, 0.35 mg/ml R-ONC. To confirm that I₁, I₂ and I₃ were produced from both N-ONC and R-ONC, co-elution tests were carried out, and the results are shown as insets in these chromatograms. Each intermediate was collected from either oxidative folding or reductive unfolding experiments, mixed together and, when re-injected into the HPLC, eluted as one peak. The absorbance at 280 nm was monitored.

unfolding pathway in ONC; this is attributed to the highly exposed (30–75) disulfide bond in N-ONC which is preferentially reduced to form I₂ (Narayan *et al.* 2004; Xu *et al.* 2004). Also, 6 M GdnHCl was required to expose the other disulfide bonds to produce I₃ in reductive unfolding.

In order to produce I₁ by reductive unfolding for trypsin digestion, the production of I₁ from I₂ is shown in Fig. 5. I₂ was used because it was shown not to reshuffle under the conditions for reduction, and the yield is much higher for I₁ than a reduction from N when compared with Fig. 4 (R.F. Gahl and H.A. Scheraga, in preparation). In addition, I₁ was produced by reducing unblocked I₃, as shown in Fig. 6. I₂ and I₃ have three disulfides each, and the reduction of these two intermediates can possibly generate three two-disulfide-bond-containing species. In Figs 5 and 6, only




Fig. 5. Chromatogram for the production of I₁ from isolated and unblocked I₂ (des-[30–75]) after 1 h. The conditions were: 25°C, 6.6 mM TCEP, pH 3, 6 M GdnHCl, 0.3 mg/ml unblocked I₂. The absorbance at 280 nm was monitored. The chromatogram shown in Fig. 1 is superimposed for comparison.

two peaks are visible between I₂ and I₃ and the reduced ONC. It is possible that two unstructured 2S species are eluted at the same time or one 2S species is not produced abundantly and thus only two peaks are detected in the chromatogram besides the original intermediate and the reduced form. The conditions to produce I₁ from I₃ (0.1 M TCEP) are different than those used to produce I₁ from I₂ (6.6 mM TCEP, 6 M GdmCl). This is a reflection of the relative reactivities around the (30–75) disulfide bond in I₃ and the (19–68) disulfide bond in I₂. The (19–68) disulfide must be buried in des-[30–75] because GdnHCl was required to expose it for reduction. Also, the (30–75) disulfide bond was exposed in des-[19–68] but is not as reactive as when it is reduced in N-ONC (Narayan *et al.* 2004; Xu *et al.* 2004). These interactions will have implications for determining the oxidative folding pathway of ONC (R.F. Gahl and H.A. Scheraga, in preparation).

I₂ and I₃ are intermediates with three disulfide bonds, whereas I₁ has two. In ‘Populating the oxidative folding intermediates by reductive unfolding for disulfide mapping’ section, it was demonstrated that one disulfide bond was reduced in I₂ to produce I₁. Since ONC is a four disulfide-bond-containing protein, and if I₁ could also be populated by another intermediate with three disulfide bonds, namely I₃, then I₁ must lack the disulfide bonds missing in both I₂ and I₃. This is indeed the case, and the results are discussed in ‘Identity of I₁, I₂ and I₃’ section.

Methodology to identify the oxidative folding intermediates, I₁, I₂ I₃

In order to identify the oxidative folding intermediates, i.e. map the disulfide linkages, two questions must be answered: (i) which particular cysteine is involved in a disulfide bond? (ii) If it is, what is its connectivity? A method developed by Watson and co-workers (Wu and Watson 1997) was adopted




Fig. 6. (A) I_3 can be reduced with 0.1 M TCEP (pH 3.1, 0.1 M citric acid, 0.1 mg/ml protein, 25°C) to form I_1 after 40 min, and separated by RP-HPLC as described in ‘Populating the oxidative folding intermediates for disulfide mapping’ section. (B) Separation of all the oxidative folding intermediates produced with 25 mM DTT^{ox} after 30 min (0.35 mg/ml protein, 25°C, pH 8.0, 0.1 M Tris) using the same RP-HPLC method. The absorbance at 210 nm was monitored.

to identify the free cysteines in I_2 and I_3 . Tryptic digestion was used to identify the free cysteines in I_1 . While each of these methods for mapping the intermediates determines whether a cysteine is involved in a disulfide bond or not, the disulfide-bond connectivity still had to be determined. To determine whether the connectivity of the disulfide bonds in the oxidative folding intermediates is native, the intermediates are also populated by *reductive unfolding* of N-ONC whose disulfide bonds are known from NMR (Gorbatyuk *et al.* 2004) and crystallography (Mosimann *et al.* 1994). The oxidative folding intermediates produced by reductive unfolding should have native disulfide bonds if they are not preceded by a species that reshuffles during the reduction. To eliminate reshuffling, which could produce non-native disulfide bonds, the reductive unfolding was performed at pH 3 with TCEP. To ensure access to all of the disulfide bonds in ONC, the reduction was performed in the presence of 6 M GdnHCl.

Identity of I_1 , I_2 and I_3

Table II shows the possible fragments that are produced using the method developed by Wu and Watson (1997). The assignment of the disulfides to I_3 is unambiguous. All three peaks in the mass spectrum corresponding to des-[19–68] are present, and the peaks corresponding to other ‘des’ species are absent. The same is true for the assignment to I_2 as des-[30–75].

The peptides of I_1 generated by trypsin that identify it as des-[19–68, 30–75] are listed in Table III. The peptides from I_1 populated by oxidative folding (I_1^{ox}) and reductive unfolding (I_1^{red}) are compared to the peptides generated by

Table II. Results of fragmentation of I_2 and I_3 using the method of Wu and Watson (1997)

I_2^{ox} ^a	I_2^{red} ^a
3516.6 ^b (1–29-Nu., 3516.6 ^c)	3516.6 (1–29-Nu., 3516.6)
5079.8 (itc-30–74-Nu., 5078.8)	5079.2 (itc-30–74-Nu., 5078.8)
3343.6 (itc-75–104, 3343.6)	3343.5 (itc-75–104, 3343.6)
I_3^{ox} ^d	I_3^{red} ^d
2241.1 (1–18-Nu., -2241.5)	2241.1 (1–18-Nu., 2241.5)
5600.0 (itc-19–67-Nu., 5596.5)	5590.8 (itc-19–67-Nu., 5596.5)
4107.4 (itc-68–104, 4103.8)	4103.2 (itc-68–104, 4103.8)

The superscripts ‘red’ and ‘ox’ refer to an intermediate produced by reductive unfolding and oxidative folding, respectively. The abbreviations –itc and –Nu: refer to the same modifications of the intermediates described in Fig. 2; these modifications are taken into account in determining the masses of the fragments. The fragments in each of the three rows identify the positions of the free cysteines. Since the same fragments are observed in fragmenting an intermediate produced by reductive unfolding or by oxidative folding and the intermediates from oxidative folding and reductive unfolding co-elute, the disulfide bonds in the intermediates are native.

^a I_2 is des-[30–75].

^bThe numbers in these columns are the masses obtained by Mass Spectroscopy.

^cThe theoretical masses for each of the fragments.

^d I_3 is des-[19–68].

RB-ONC under the same conditions, and the two disulfide bonds were identified as Cys87–Cys104 and Cys48–Cys90.

The corresponding mass spectra to produce Tables II and III are provided in the Supplementary Material (Supplementary data are available at PEDS online).

Table III. Results from tryptic digestion of RB-ONC, I₁^{ox}, and I₁^{red}

Mass of peptides	Location of peptides in ONC	Cys state	RB-ONC	I ₁ ^{red} ^a	I ₁ ^{ox} ^a
1994.9 (1994.9)	86–104	C87-C104, C90-SH	No	Yes	Yes
2004.8 (2004.8)	16–31	C19blo, C30blo	Yes	No	No
2221.9 (2221.9)	86–104	C87blo, C90blo, C104blo	Yes	No	No
2426.1 (2426.1)	46–49...86–104	C48-C90, C87-C104	No	Yes	Yes
2527.1 (2527.1)	56–76	C68blo, C75blo	Yes	No	No
5561.8 (5562.1)	1–45	C19blo, C30blo	Yes	Yes	Yes

The superscript ‘red’ refers to the population produced by reductive unfolding, and the superscript ‘ox’ refers to the population produced by oxidative folding. Cysteines in a fragment blocked with an $-\text{SCH}_2\text{CH}_2\text{NH}_3^+$ group from AEMTS are indicated as CXXblo, or in a disulfide bond as CXX-CXX, and unblocked as CXX-SH. The numbers in the first part of column 1 are the experimental masses, and those in parentheses are the theoretical masses of the fragments. Since the same fragments are produced from AEMTS-blocked I₁ populated by either reductive unfolding or oxidative folding, I₁ must have native disulfide bonds. ‘Yes’ and ‘No’ indicate that the particular peptide is, or is not, present in the mass spectrum.

^aI₁ is des-[19–68, 30–75].

Discussion

The oxidative folding of ONC with DTT^{ox} has been examined at 25°C (pH 8). Three peaks that contain intermediates with disulfide bonds (I₁, I₂ and I₃) are found to survive a reduction pulse, indicating that the intermediates constituting these peaks have some degree of structure that protects their disulfides from a reducing agent. Further studies (in preparation), including how they are involved in the oxidative folding pathway, will provide insight into the process by which ONC obtains its native structure from its completely reduced form. Preliminary oxidative folding data indicate that I₁ is formed and consumed early in the regeneration process. While I₂ and I₃ are produced after I₁ first appears, I₂ is consumed more quickly before the end of the regeneration; I₃ reacts more slowly relative to I₂ and, therefore, is not as reactive and not readily oxidized to N.

The formation of such a structured intermediate (I₁) in the early stages of ONC folding is compatible with the enhanced rate of regeneration. Once a structured intermediate is formed, its disulfides are sequestered from reduction and reshuffling reactions by its remaining cysteines or by an internal redox agent. Additionally, the remaining cysteines are aligned in native fashion and facilitate formation of the native disulfide bonds (Laity *et al.* 1997). Thus, I₁ would be ideally poised to be oxidized to a structured three-disulfide-containing intermediate that can then be oxidized easily to N (2S*→3S*→N). The current work is being devoted to determine how these reactions fit into an overall oxidative folding mechanism.

Although RNase A and ONC are structurally very similar, our results indicate that there are significant differences in their oxidative folding pathways. The oxidative folding of RNase A at 25°C is characterized by the presence of two stable, three-disulfide-containing on-pathway intermediates, namely des-[40–95] and des-[65–72] (Rothwarf *et al.* 1998a). Furthermore, no stable two-disulfide intermediate(s) have been found to exist in the RNase A folding pathway whereas, in ONC, the presence of a stable, two-disulfide-containing intermediate (I₁) is clearly evident. It is clear that the nature of the intermediates produced in these regenerative oxidative pathways is very different. This indicates that the intramolecular interactions that stabilize their respective intermediates during the oxidative folding

process differ, and yet are capable of giving rise to a very similar three-dimensional fold in the final stage of the folding process. By identifying the interactions that give rise to the different folding pathways in future studies, more information can be gained about the interactions that drive the folding process.

Finally, the oxidative folding of ONC is of special interest because it lacks the (65–72) disulfide bond that plays a major role in the oxidative folding of RNase A. In the unstructured intermediates of RNase A, this bond is present to an extent of ~40% of all the disulfide bonds in the 1S and 2S ensembles (Xu *et al.* 1996; Volles *et al.* 1999), and the structured intermediate lacking this disulfide bond (des-[65–72]) is formed to a minor extent (~20%) in the rate-determining step of its regeneration (Rothwarf *et al.* 1998b). It is, therefore, of interest that the oxidative folding of ONC is faster than that of RNase A, despite the absence of the analog of the (65–72) disulfide bond in ONC. Such a disulfide bond analog does not appear in the native structure of ONC. ONC has two cysteines, Cys68 and Cys75, the same distance apart along the peptide backbone as the (65–72) disulfide bond in RNase A. However, in N-ONC, Cys68 is involved in a disulfide bond with Cys19, and Cys75 is involved in a disulfide bond with Cys30. Each of these disulfide bonds are 39 and 45 residues apart, respectively, and their formation incurs a greater loss in entropy for the polypeptide than forming a disulfide bond seven residues apart. However, the formation of the 19–68 and 30–75 disulfide bonds may be driven by enthalpically favorable interactions of the side chains. A similar preference to form a disulfide by enthalpic interactions is seen in the case of RNase A. Cys58 and Cys65 are seven residues apart, but Cys65 forms a disulfide bond with Cys72 because the (65–72) bond is a native disulfide bond, and more favorable enthalpic interactions promote this disulfide preference. Cys58 forms a native disulfide bond with Cys110.

In conclusion, we have demonstrated that the oxidative folding pathway of ONC is significantly different from that of its structural homologue, RNase A. Early-forming, structured intermediates can significantly accelerate the oxidative folding rate. Further experiments are in progress to identify the intramolecular interactions that account for the increased rate and formation of the structured intermediates.

Acknowledgements

We thank Dr. Ervin Welker for assistance in the expression of ONC and for helpful comments during the course of this work and also thank Robert Sherwood at the Cornell University Life Sciences Core Laboratories Center Proteomics and Mass Spectrometry Facility for obtaining the mass spectra.

Funding

National Institute of General Medical Sciences of the National Institutes of Health (Grant no. GM-24893).

References

- Ardelt,W., Mikulski,S.M. and Shogen,K. (1991) *J. Biol. Chem.*, **266**, 245–251.
- Boix,E., Wu,Y., Vasandani,V.M., Saxena,S.K., Ardel,W., Ladner,J. and Youle,R.J. (1996) *J. Mol. Biol.*, **257**, 992–1007.
- Chang,J.Y. and Li,L. (2002) *Biochemistry*, **41**, 8405–8413.
- Gorbatyuk,V.Y., Tsai,C.-K., Chang,C.-F. and Huang,T. (2004) *J. Biol. Chem.*, **279**, 5772–5780.
- Iwaoka,M., Juminaga,D. and Scheraga,H.A. (1998) *Biochemistry*, **37**, 4490–4501.
- Laity,J.H., Shimotakahara,S. and Scheraga,H.A. (1993) *Proc. Natl Acad. Sci. USA*, **90**, 615–619.
- Laity,J.H., Lester,C.C., Shimotakahara,S., Zimmerman,D.E., Montelione,G.T. and Scheraga,H.A. (1997) *Biochemistry*, **36**, 12683–12699.
- Leland,P.A. and Raines,R.T. (2001) *Chem. Biol.*, **8**, 405–413.
- Leland,P.A., Schultz,L.W., Kim,B.M. and Raines,R.T. (1998) *Proc. Natl Acad. Sci. USA*, **95**, 10407–10412.
- Leland,P.A., Staniszewski,K.E., Kim,B.M. and Raines,R.T. (2000) *FEBS Lett.*, **477**, 203–207.
- Mosimann,S.C., Ardel,W. and James,M.N.G. (1994) *J. Mol. Biol.*, **236**, 1141–1153.
- Narayan,M., Welker,E., Wedemeyer,W.J. and Scheraga,H.A. (2000) *Acc. Chem. Res.*, **33**, 805–812.
- Narayan,M., *et al.* (2004) *J. Mol. Biol.*, **338**, 795–809.
- Newton,D.L., Boque,L., Wlodawer,A., Huang,C.Y. and Rybak,S.M. (1998) *Biochemistry*, **37**, 5173–5183.
- Newton,D.L., Hansen,H.J., Mikulski,S.M., Goldenberg,D.M. and Rybak,S.M. (2001) *Blood*, **97**, 528–535.
- Rothwarf,D.M., Li,Y.J. and Scheraga,H.A. (1998a) *Biochemistry*, **37**, 3760–3766.
- Rothwarf,D.M., Li,Y.J. and Scheraga,H.A. (1998b) *Biochemistry*, **37**, 3767–3776.
- Rothwarf,D.M. and Scheraga,H.A. (1993) *Biochemistry*, **32**, 2671–2679.
- Saxena,S.K., Sirdeshmukh,R., Ardel,W., Mikulski,S.M., Shogen,K. and Youle,R.J. (2002) *J. Biol. Chem.*, **277**, 15142–15146.
- Thannhauser,T.W., Rothwarf,D.M. and Scheraga,H.A. (1997) *Biochemistry*, **36**, 2154–2165.
- Volles,M.J., Xu,X. and Scheraga,H.A. (1999) *Biochemistry*, **38**, 7284–7293.
- van den Berg,B., Chung,E.W., Robinson,C.V., Mateo,P.L. and Dobson,C.M. (1999) *EMBO J.*, **18**, 4794–4803.
- Wedemeyer,W.J., Welker,E., Narayan,M. and Scheraga,H.A. (2000) *Biochemistry*, **39**, 4207–4216.
- Weissman,J.S. and Kim,P.S. (1991) *Science*, **253**, 1386–1393.
- Welker,E., Narayan,M., Wedemeyer,W.J. and Scheraga,H.A. (2001a) *Proc. Natl Acad. Sci. USA*, **98**, 2312–2316.
- Welker,E., Wedemeyer,W.J., Narayan,M. and Scheraga,H.A. (2001b) *Biochemistry*, **40**, 9059–9064.
- Welker,E., Hathaway,L. and Scheraga,H.A. (2004) *J. Am. Chem. Soc.*, **126**, 3270–3271.
- Welker,E., Hathaway,L., Xu,G., Narayan,M., Pradeep,L., Shin,H.-C. and Scheraga,H.A. (2007) *Biochemistry*, **46**, 5485–5493.
- Wlodawer,A., Svensson,L.A., Sjölin,L. and Gilliland,G.L. (1988) *Biochemistry*, **27**, 2705–2717.
- Wu,J. and Watson,J.T. (1997) *Prot. Sci.*, **6**, 391–398.
- Wu,J. and Watson,J.T. (1998) *Anal. Biochem.*, **258**, 268–276.
- Xu,X. and Scheraga,H.A. (1998) *Biochemistry*, **37**, 7561–7571.

Xu,X., Rothwarf,D.M. and Scheraga,H.A. (1996) *Biochemistry*, **35**, 6406–6417.

Xu,G., Narayan,M., Welker,E. and Scheraga,H.A. (2004) *Biochemistry*, **43**, 3246–3254.

Received November 14, 2007; revised November 14, 2007;
accepted November 20, 2007

Edited by Valerie Daggett